

Component sizes of the random graph outside the scaling window

ASAF NACHMIAS AND YUVAL PERES*

February 2, 2008

Abstract

We provide simple proofs describing the behavior of the largest component of the Erdős-Rényi random graph $G(n, p)$ outside of the scaling window, $p = \frac{1+\epsilon(n)}{n}$ where $\epsilon(n) \rightarrow 0$ but $\epsilon(n)n^{1/3} \rightarrow \infty$.

1 Introduction

Consider the random graph $G(n, p)$ obtained from the complete graph on n vertices by retaining each edge with probability p and deleting each edge with probability $1 - p$. We denote by $\mathcal{C}_j$ the j -th largest component. Let $\epsilon(n)$ be a non-negative sequence such that $\epsilon(n) \rightarrow 0$ and $\epsilon(n)n^{1/3} \rightarrow \infty$. The following theorems, proved by Bollobás [4] and Łuczak [8] using different methods, describe the behavior of the largest components when p is outside the “scaling-window”.

Theorem 1 [Subcritical phase] *If $p(n) = \frac{1-\epsilon(n)}{n}$ then for any $\eta > 0$ and integer $\ell > 0$ we have*

$$\mathbf{P}\left(\left|\frac{|\mathcal{C}_\ell|}{2\epsilon(n)^{-2}\log(n\epsilon(n)^3)} - 1\right| > \eta\right) \rightarrow 0,$$

as $n \rightarrow \infty$.

Theorem 2 [Supercritical phase] *If $p(n) = \frac{1+\epsilon(n)}{n}$ then for any $\eta > 0$ we have*

$$\mathbf{P}\left(\left|\frac{|\mathcal{C}_1|}{2n\epsilon(n)} - 1\right| > \eta\right) \rightarrow 0,$$

*Microsoft Research and U.C. Berkeley. Research of both authors supported in part by NSF grants #DMS-0244479 and #DMS-0104073

and for any integer $\ell > 1$ we have

$$\mathbf{P}\left(\left|\frac{|\mathcal{C}_\ell|}{2\epsilon^{-2}(n)\log(n\epsilon^3)} - 1\right| > \eta\right) \rightarrow 0,$$

as $n \rightarrow \infty$.

The proofs of these theorems in [4] and [8] are quite involved and use the detailed asymptotics from [14], [4] and [3] for the number of graphs on k vertices with $k + \ell$ edges. The proofs we present here are simple and require no hard theorems. The main advantage, however, of these proofs is their robustness. In a companion paper [12] we use similar methods to analyze critical percolation on a random regular graphs. In this case, the enumerative methods employed in [4] and [8] are not available.

The phase transition in the Erdős-Rényi random graphs $G(n, p)$ happens when $p = \frac{c}{n}$. Namely, with high probability, if $c > 1$ then $|\mathcal{C}_1|$ is linear in n , and if $c < 1$ then $|\mathcal{C}_1|$ is logarithmic in n . When $c \sim 1$ the situation is more delicate. In [9], Łuczak, Pittel and Wierman prove that for $p = \frac{1+\lambda n^{-1/3}}{n}$, the law of $n^{-2/3}|\mathcal{C}_1|$ converges to a positive non-constant distribution which in [1] is identified as the longest excursion length of some Brownian motion with variable drift. See [11] for a recent account of the case $p = \frac{1+\lambda n^{-1/3}}{n}$ with simple proofs.

Thus, $|\mathcal{C}_1|$ is not concentrated and is roughly of size $n^{2/3}$ if $p = \frac{1+\lambda n^{-1/3}}{n}$. However, if $\epsilon(n)$ a sequence such that $n^{1/3}\epsilon(n) \rightarrow \infty$ and $p = \frac{1+\epsilon(n)}{n}$ then as stated in Theorems 1 and 2, the size $|\mathcal{C}_1|$ of the largest component in $G(n, p)$ is concentrated. In summary, $G(n, p)$ has a scaling window of length $n^{-1/3}$ in which the percolation is “critical” in the sense that $|\mathcal{C}_1|$ is not concentrated.

2 The exploration process

We recall an exploration process, due to Karp and Martin-Löf (see [7] and [10]), in which vertices will be either *active*, *explored* or *neutral*. After the completion of step $t \in \{0, 1, \dots, n\}$ we will have precisely t explored vertices and the number of the active and neutral vertices is denoted by A_t and N_t respectively.

Fix an ordering of the vertices $\{v_1, \dots, v_n\}$. In step $t = 0$ of the process, we declare vertex v_1 *active* and all other vertices *neutral*. Thus $A_0 = 1$ and $N_0 = n - 1$. In step $t \in \{1, \dots, n\}$, if $A_{t-1} > 0$ let w_t be the first active vertex; if $A_{t-1} = 0$, let w_t be the first neutral vertex. Denote by η_t the

number of neutral neighbors of w_t in $G(n, p)$, and change the status of these vertices to *active*. Then, set w_t itself *explored*.

Denote by $\mathcal{F}_t$ the σ -algebra generated by $\{\eta_1, \dots, \eta_t\}$. Observe that given $\mathcal{F}_{t-1}$ the random variable η_t is distributed as $\text{Bin}(N_{t-1} - \mathbf{1}_{\{A_{t-1}=0\}}, p)$ and we have the recursions

$$N_t = N_{t-1} - \eta_t - \mathbf{1}_{\{A_{t-1}=0\}}, \quad t \leq n, \quad (1)$$

and

$$A_t = \begin{cases} A_{t-1} + \eta_t - 1, & A_{t-1} > 0 \\ \eta_t, & A_{t-1} = 0, \end{cases} \quad t \leq n. \quad (2)$$

As every vertex is either neutral, active or explored,

$$N_t = n - t - A_t, \quad t \leq n. \quad (3)$$

At each time $j \leq n$ in which $A_j = 0$, we have finished exploring a connected component. Hence the random variable Z_t defined by

$$Z_t = \sum_{j=1}^{t-1} \mathbf{1}_{\{A_j=0\}},$$

counts the number of components completely explored by the process before time t . Define the process $\{Y_t\}$ by $Y_0 = 1$ and

$$Y_t = Y_{t-1} + \eta_t - 1.$$

By (2) we have that $Y_t = A_t - Z_t$, i.e. Y_t counts the number of active vertices at step t minus the number of components completely explored before step t .

At each step we marked as explored precisely one vertex. Hence, the component of v_1 has size $\min\{t \geq 1 : A_t = 0\}$. Moreover, let $t_1 < t_2 \dots$ be the times at which $A_{t_j} = 0$; then $(t_1, t_2 - t_1, t_3 - t_2, \dots)$ are the sizes of the components. Observe that $Z_t = Z_{t_j} + 1$ for all $t \in \{t_j + 1, \dots, t_{j+1}\}$. Thus $Y_{t_{j+1}} = Y_{t_j} - 1$ and if $t \in \{t_j + 1, \dots, t_{j+1} - 1\}$ then $A_t > 0$, and thus $Y_{t_{j+1}} < Y_t$. By induction we conclude that $A_t = 0$ if and only if $Y_t < Y_s$ for all $s < t$, i.e. $A_t = 0$ if and only if $\{Y_t\}$ has hit a new record minimum at time t . By induction we also observe that $Y_{t_j} = -(j - 1)$ and that for $t \in \{t_j + 1, \dots, t_{j+1}\}$ we have $Z_t = j$. Also, by our previous discussion for $t \in \{t_j + 1, \dots, t_{j+1}\}$ we have $\min_{s \leq t-1} Y_t = Y_{t_j} = -(j - 1)$, hence by induction we deduce that $Z_t = -\min_{s \leq t-1} Y_t + 1$. Consequently,

$$A_t = Y_t - \min_{s \leq t-1} Y_s + 1. \quad (4)$$

Lemma 3 *For all $p \leq \frac{2}{n}$ there exists a constant $c > 0$ such that for any integer $t > 0$,*

$$\mathbf{P}\left(N_t \leq n - 5t\right) \leq e^{-ct}.$$

PROOF. Let $\{\alpha_i\}_{i=1}^t$ be a sequence of i.i.d. random variables distributed as $\text{Bin}(n, p)$. It is clear that we can couple η_i and α_i so $\eta_i \leq \alpha_i$ for all i , and thus by (1)

$$N_t \geq n - 1 - t - \sum_{i=1}^t \alpha_i. \quad (5)$$

The sum $\sum_{i=1}^t \alpha_i$ is distributed as $\text{Bin}(nt, p)$ and $p \leq \frac{2}{n}$ so by Large Deviations (see [2] section A.14) we get that for some fixed $c > 0$

$$\mathbf{P}\left(\sum_{i=1}^t \alpha_i \geq 3t\right) \leq e^{-ct},$$

which together with (5) concludes the proof. $\square$

3 The subcritical phase

Before beginning the proof of Theorem 1 we require some facts about processes with i.i.d. increments. Fix some small $\epsilon > 0$ and let $p = \frac{1-\epsilon}{m}$ for some integer $m > 1$. Let $\{\beta_j\}$ be a sequence of random variables distributed as $\text{Bin}(m, p)$. Let $\{W_t\}_{t \geq 0}$ be a process defined by

$$W_0 = 1, \quad W_t = W_{t-1} + \beta_t - 1.$$

Let τ be the hitting time of 0,

$$\tau = \min_t \{W_t = 0\}.$$

By Wald's lemma we have that $\mathbf{E} \tau = \epsilon^{-1}$. Further information on the tail distribution of τ is given by the following lemma.

Lemma 4 *There exists constant $C_1, C_2, c_1, c_2 > 0$ such that for all $T > \epsilon^{-2}$ we have*

$$\mathbf{P}(\tau \geq T) \leq C_1 \left(\epsilon^{-2} T^{-3/2} e^{-\frac{(\epsilon^2 - c_1 \epsilon^3)T}{2}} \right),$$

and

$$\mathbf{P}(\tau \geq T) \geq c_1 \left(\epsilon^{-2} T^{-3/2} e^{-\frac{(\epsilon^2 + c_2 \epsilon^3)T}{2}} \right).$$

Furthermore,

$$\mathbf{E} \tau^2 = O(\epsilon^{-3}).$$

We will use the following proposition due to Spitzer (see [13]).

Proposition 5 *Let $a_0, \dots, a_{k-1} \in \mathbb{Z}$ satisfy $\sum_{i=0}^{k-1} a_i = -1$. Then there is precisely one $j \in \{0, \dots, k-1\}$ such that for all $r \in \{0, \dots, k-2\}$*

$$\sum_{i=0}^r a_{(j+i) \bmod k} \geq 0.$$

Proof of Lemma 4. By Proposition 5, $\mathbf{P}(\tau = t) = \frac{1}{t} \mathbf{P}(W_t = 0)$. As $\sum_{j=1}^t \beta_j$ is distributed as a $\text{Bin}(mt, p)$ random variable we have

$$\mathbf{P}(W_t = 0) = \binom{mt}{t-1} p^{t-1} (1-p)^{m-(t-1)}.$$

Replacing $t-1$ with t in the above formula only changes it by a multiplicative constant which is always between $1/2$ and 2 . A straightforward computation using Stirling's approximation gives

$$\mathbf{P}(W_t = 0) = \Theta \left\{ t^{-1/2} (1-\epsilon)^t \left(1 + \frac{1}{m-1}\right)^{tm} \left(1 - \frac{1-\epsilon}{m}\right)^{t(m-1)} \right\}. \quad (6)$$

Denote $q = (1-\epsilon) \left(1 + \frac{1}{m-1}\right)^m \left(1 - \frac{1-\epsilon}{m}\right)^{m-1}$, then

$$\mathbf{P}(\tau \geq T) = \sum_{t \geq T} \mathbf{P}(\tau = t) = \sum_{t \geq T} \frac{1}{t} \mathbf{P}(W_t = 0) = \Theta \left(\sum_{t \geq T} t^{-3/2} q^t \right).$$

This sum can be bounded above by

$$T^{-3/2} \sum_{t \geq T} q^t = T^{-3/2} \frac{q^T}{1-q},$$

and below by

$$\sum_{t=T}^{2T} t^{-3/2} q^t \geq (2T)^{-3/2} \frac{q^T (1-q^T)}{1-q}.$$

Observe that as $m \rightarrow \infty$ we have that q tends to $(1-\epsilon)e^\epsilon$. By expanding e^ϵ we find that

$$q = (1-\epsilon) \left(1 + \epsilon + \frac{\epsilon^2}{2}\right) + \Theta(\epsilon^3) = 1 - \frac{\epsilon^2}{2} + \Theta(\epsilon^3).$$

Using this and the previous bounds on $\mathbf{P}(\tau \geq T)$ we get the first assertion of the Lemma.

The second assertion follows from the following computation. By (6) we have that for some constant $C > 0$

$$\mathbf{E} \tau^2 = \sum_{t \geq 1} t^2 \mathbf{P}(\tau = t) = \sum_{t \geq 1} t \mathbf{P}(W_t = 0) \leq C \sum_{t \geq 1} \sqrt{t} q^t.$$

Thus, by direct computation (or by [6], section XIII.5, Theorem 5)

$$\mathbf{E} \tau^2 \leq O\left(\frac{1}{1-q}\right)^{3/2} = O(\epsilon^{-3}).$$

□

Proof of Theorem 1. We begin with an upper bound. Recall that component sizes are $t_{j+1} - t_j$ for some $j > 0$ where t_j are record minima of the process $\{Y_t\}$. For a vertex v denote by $C(v)$ the connected component of $G(n, p)$ which contains v . We first bound $\mathbf{P}(|C(v_1)| > T)$ where

$$T = 2(1 + \eta)\epsilon^{-2} \log(n\epsilon^3).$$

Recall that $|C(v_1)| = \min_t \{Y_t = 0\}$. Couple $\{Y_t\}$ with a process $\{W_t\}$ as in Lemma 4, which has increments distributed as $\text{Bin}(n, p) - 1$ such that $Y_t \leq W_t$ for all t . Define τ as in Lemma 4. As $p = \frac{1-\epsilon}{n}$ and $T > \epsilon^{-2}$, by Lemma 4 we have

$$\mathbf{P}(\tau > T) \leq C\epsilon(n\epsilon^3)^{-(1+\eta)} \log(n\epsilon^3)^{-3/2},$$

for some fixed $C > 0$. Our coupling implies that $\mathbf{P}(|C(v_1)| > T) \leq \mathbf{P}(\tau > T)$. Denote by X the number of vertices v such that $|C(v)| > T$. If $|\mathcal{C}_1| > T$ then $X > T$. Also, for any two vertices v and u by symmetry we have that $|C(v)|$ and $|C(u)|$ are identically distributed. We conclude that

$$\begin{aligned} \mathbf{P}(|\mathcal{C}_1| > T) &\leq \mathbf{P}(X > T) \leq \frac{\mathbf{E} X}{T} = \frac{n\mathbf{P}(|C(v_1)| > T)}{T} \\ &\leq \frac{C_1 n \epsilon (n\epsilon^3)^{-(1+\eta)(1-C_2\epsilon)} \log^{-3/2}(n\epsilon^3)}{2(1 + \eta)\epsilon^{-2} \log(n\epsilon^3)} \leq (n\epsilon^3)^{-\eta(1-C_2\epsilon)+C_2\epsilon} \rightarrow 0. \end{aligned}$$

We now turn to prove a lower bound. Write

$$T = 2(1 - \eta)\epsilon^{-2} \log(n\epsilon^3),$$

and define the stopping time

$$\gamma = \min\{t : N_t \leq n - \frac{\eta\epsilon n}{8}\}.$$

Recall that $\{t_j\}$ are times in which $A_{t_j} = 0$ and also Y_{t_j} is a record minimum for $\{Y_t\}$. For each integer j let $\{W_t^{(j)}\}$ be a process with increments distributed as $\text{Bin}(n - \frac{\eta\epsilon n}{8}, p)$ where the starting point is $W_0^{(j)} = Y_{t_j} = -(j-1)$. Note that if $t_{j+1} < \gamma$ then we can couple $\{Y_t\}$ and $\{W_t^{(j)}\}$ such that $Y_{t_j+t} \geq W_t$ for all $t \in [t_j, t_{j+1}]$. Define the stopping times $\{\tau_j\}$ by

$$\tau_j = \min\{t : W_t^{(j)} = -j\}.$$

Take

$$N = \left\lceil \epsilon^{-1}(n\epsilon^3)^{(1-\frac{\eta}{8})} \right\rceil.$$

We will prove that with high probability $t_N < \gamma$ and that there exists $k_1 < k_2 < \dots < k_\ell < N$ such that $\tau_{k_i} > T$. Note that these two events imply that $|\mathcal{C}_\ell| > T$. Indeed, by Lemma 3 we have

$$\mathbf{P}\left(\gamma \leq \frac{\eta\epsilon n}{40}\right) \leq e^{-c\epsilon n}. \quad (7)$$

By bounding the increments of $\{Y_t\}$ above by variables distributed as $\text{Bin}(n, p) - 1$ we learn by Wald's Lemma (see [5]) that $\mathbf{E}[t_{j+1} - t_j] \leq \epsilon^{-1}$, hence $\mathbf{E}t_N \leq \epsilon^{-2}(n\epsilon^3)^{(1-\frac{\eta}{8})}$. We conclude that

$$\mathbf{P}(t_N > \frac{\eta\epsilon n}{40}) \leq \frac{40\epsilon^{-2}(n\epsilon^3)^{(1-\frac{\eta}{8})}}{\eta\epsilon n} = \frac{40}{\eta}(n\epsilon^3)^{-\frac{\eta}{8}}, \quad (8)$$

which goes to 0 as $\epsilon n^{-1/3}$ tends to ∞ . In Lemma 4 take $m = n - \frac{\eta\epsilon n}{8}$ and note that $p = \frac{(1-\epsilon)(1-\frac{\eta\epsilon}{8})}{m} \geq \frac{1-(1+\frac{\eta}{8})\epsilon}{m}$, and so Lemma 4 gives that for any j

$$\mathbf{P}(\tau_j > T) \geq c_1\epsilon(n\epsilon^3)^{-(1+\frac{\eta}{8})^2(1-\eta)(1+c_2\epsilon)} \log^{-3/2}(\epsilon^3 n) \geq \epsilon(n\epsilon^3)^{-(1-\frac{\eta}{4})}.$$

Let X be the number of $j \leq N$ such that $\tau_j > T$. Then we have

$$\mathbf{E}X \geq N\epsilon(n\epsilon^3)^{-(1-\frac{\eta}{4})} \geq C(n\epsilon^3)^{\frac{\eta}{8}} \rightarrow \infty,$$

hence by Large Deviations (see [2], section A.14) for any fixed integer $\ell > 0$ we have

$$\mathbf{P}(X < \ell) \leq e^{-c(n\epsilon^3)^{\frac{\eta}{8}}},$$

for some fixed $c > 0$. By our previous discussion, this together with (7) and (8) gives

$$\mathbf{P}(|\mathcal{C}_\ell| < T) \leq O\left(\frac{(n\epsilon^3)^{-\frac{\eta}{8}}}{\eta}\right).$$

□

4 The supercritical phase

In this section we denote $\xi_t = \eta_t - 1$. We first prove some Lemmas.

Lemma 6 *If $p = \frac{1+\epsilon}{n}$ then for all $t < 3\epsilon(n)n$*

$$\mathbf{E} A_t = O(\epsilon t + \sqrt{t}), \quad (9)$$

and

$$\mathbf{E} Z_t = O(\epsilon t + \sqrt{t}). \quad (10)$$

PROOF. Write $T = 3\epsilon n$. We will use (4). First observe that as η_t can always be bounded above by a $\text{Bin}(n, p)$ random variable we can bound $\mathbf{E} \xi_t \leq \epsilon$ for all t and hence $\mathbf{E} Y_t \leq \epsilon t$. Denote by τ the stopping time $\tau = \min\{t : N_t \leq n - 15\epsilon n\}$. By definition of η_t we have

$$\mathbf{E} [\xi_t \mid \mathcal{F}_{t-1}] = pN_{t-1} - p\mathbf{1}_{\{A_{t-1}=0\}} - 1.$$

As $\{N_t\}$ is a decreasing sequence, we deduce that as long as $t < \tau$, we have $\mathbf{E} [\xi_t \mid \mathcal{F}_{t-1}] > -D\epsilon$ for $D > 0$ large enough. Hence, the process $\{D\epsilon j - Y_j\}_{j=0}^{t \wedge \tau}$ is a submartingale for any t . By Doob's maximal L^2 inequality we have

$$\mathbf{E} [\max_{j \leq t \wedge \tau} (D\epsilon j - Y_j)^2] \leq 4\mathbf{E} [(D\epsilon(t \wedge \tau) - Y_{t \wedge \tau})^2]. \quad (11)$$

For any $j < \tau$ the random variable η_j can be stochastically bounded from below by a $\text{Bin}(n - 15\epsilon n, p)$ random variable and above by a $\text{Bin}(n, p)$ random variable. Hence for any $k < j < \tau$ we have

$$\left| \mathbf{E} [\xi_j - D\epsilon \mid \mathcal{F}_k] \right| = O(\epsilon),$$

and so

$$\mathbf{E} [(\xi_j - D\epsilon)(\xi_k - D\epsilon)] = O(\epsilon^2).$$

We conclude that as long as $t < \tau$

$$\mathbf{E}[(D\epsilon t - Y_t)^2] \leq 2 \sum_{k < j}^t \mathbf{E}[(\xi_j - D\epsilon)(\xi_k - D\epsilon)] + \sum_{j=1}^t \mathbf{E}[(\xi_j - D\epsilon)^2] = O(\epsilon^2 t^2 + t).$$

Lemma 3 implies that for n large enough,

$$\mathbf{P}(N_T \leq n - 15\epsilon n) \leq e^{-3c\epsilon n} \leq \frac{1}{n^2}, \quad (12)$$

and as $\{N_t\}$ is a decreasing sequence we deduce that $\mathbf{P}(\tau \leq T) \leq n^{-2}$. Hence for any $t \leq T$

$$\begin{aligned} \mathbf{E}[(D\epsilon t - Y_t)^2] &\leq \mathbf{E}[(D\epsilon(t \wedge \tau) - Y_{t \wedge \tau})^2 \mathbf{1}_{\{t < \tau\}}] + O(n^2) \mathbf{P}(t \geq \tau) \\ &= O(\epsilon^2 t^2 + t). \end{aligned}$$

We deduce by (11) and Jensen inequality that for any $t \leq T$

$$\mathbf{E}[\min_{j \leq t} (Y_j - D\epsilon j)] = O(\epsilon t + \sqrt{t}),$$

hence $\mathbf{E}[\min_{j \leq t} Y_j] = O(\epsilon t + \sqrt{t})$ and so by (4) we obtain (9). Inequality (10) follows immediately from the relation $Z_t = A_t - Y_t$. $\square$

Lemma 7 *If $p = \frac{1+\epsilon}{n}$ then for all $t < 3\epsilon(n)n$*

$$\mathbf{E} N_t = n(1-p)^t + O(\epsilon^2 n), \quad (13)$$

and

$$\mathbf{E} \xi_t = \epsilon - \frac{t}{n} + O(\epsilon^2). \quad (14)$$

PROOF. Observe that by (1) we have that

$$\mathbf{E}[N_t \mid \mathcal{F}_{t-1}] = (1-p)N_{t-1} - (1-p)\mathbf{1}_{\{A_{t-1}=0\}}.$$

By iterating this relation we get that $\mathbf{E} N_t = n(1-p)^t + O(\mathbf{E} Z_t)$ which by Lemma 6 yields (13) (observe that for $t = 3\epsilon n$ we have $\epsilon t > \sqrt{t}$ by our assumption on ϵ). Since

$$E[\xi_t \mid \mathcal{F}_{t-1}] = pN_{t-1} - p\mathbf{1}_{\{A_{t-1}=0\}} - 1,$$

by taking expectations and using (13) we get

$$\begin{aligned}
\mathbf{E} \xi_t &= (1 + \epsilon) \left(1 - \frac{1 + \epsilon}{n}\right)^t - 1 + O(\epsilon^2) \\
&= (1 + \epsilon) \left(1 - (1 + \epsilon)t/n\right) - 1 + O(\epsilon^2) = \epsilon - \frac{t}{n} + O(\epsilon^2),
\end{aligned}$$

where we used the fact that $(1 - x)^t = 1 - tx + O(t^2 x^2)$. $\square$

Proof of Theorem 2. Write $T = 3\epsilon n$ and $\xi_j^* = \mathbf{E}[\xi_j \mid \mathcal{F}_{j-1}]$. The process

$$M_t = Y_t - \sum_{j=1}^t \xi_j^*,$$

is a martingale. By Doob's maximal L^2 inequality we have that

$$\mathbf{E}(\max_{t \leq T} M_t^2) \leq 4\mathbf{E} M_T^2.$$

As M_t has orthogonal increments with bounded second moment we conclude that $\mathbf{E} M_T^2 = O(T)$, hence, by Jensen's inequality we have

$$\mathbf{E} \left[\max_{t \leq T} \left| Y_t - \sum_{j=1}^t \xi_j^* \right| \right] \leq O(\sqrt{T}) = O(\sqrt{\epsilon n}). \quad (15)$$

As $\xi_j^* = pN_{j-1} - p\mathbf{1}_{\{A_{j-1}=0\}} - 1$ by (3) we have

$$\mathbf{E} |\xi_j^* - \mathbf{E} \xi_j| = p\mathbf{E} |A_{j-1} + \mathbf{1}_{\{A_{j-1}=0\}} - \mathbf{E} A_{j-1} - \mathbf{E} \mathbf{1}_{\{A_{j-1}=0\}}|.$$

By the triangle inequality and Lemma 6 we conclude that for all $j \leq T$

$$\mathbf{E} |\xi_j^* - \mathbf{E} \xi_j| \leq p \cdot O(\epsilon j + \sqrt{j}),$$

and hence for any $t \leq T$

$$\mathbf{E} \left[\sum_{j \leq t} |\xi_j^* - \mathbf{E} \xi_j| \right] \leq p \cdot O(\epsilon t^2 + t^{3/2}) \leq O(\epsilon^3 n).$$

By the triangle inequality we get

$$\mathbf{E} \left[\max_{t \leq T} \left| \sum_{j=1}^t (\xi_j^* - \mathbf{E} \xi_j) \right| \right] \leq O(\epsilon^3 n). \quad (16)$$

Using the triangle inequality, (15), (16) and Markov inequality gives

$$\mathbf{P}\left(\max_{t \leq T} \left|Y_t - \sum_{j=1}^t \mathbf{E} \xi_j\right| \geq \delta \epsilon^2 n\right) \leq \delta^{-1}(O(\epsilon) + O((\epsilon^3 n)^{-1/2})) \longrightarrow 0. \quad (17)$$

Lemma 7 implies that for any $b > 0$

$$\sum_{j=1}^{b\epsilon n} \mathbf{E} \xi_j = \sum_{j=1}^{b\epsilon n} \left(\epsilon - \frac{t}{n} + O(\epsilon^2)\right) = (b - \frac{b^2}{2})\epsilon^2 n + O(\epsilon^3 n). \quad (18)$$

By (17) and (18) we deduce that for $\eta > 0$ small enough, with probability tending to 1, the process Y_t is strictly positive at times $[\eta\epsilon n, (2 - \eta)\epsilon n]$ and hence

$$\mathbf{P}\left(|\mathcal{C}_1| > 2(1 - \eta)\epsilon n\right) \geq 1 - O\left(\eta^{-1}(\epsilon + (\epsilon^3 n)^{-1/2})\right).$$

We also deduce by (17) and (18) that at time $t = (2 + \eta)\epsilon n$ we have $Y_t \leq -\frac{\eta^2}{3}\epsilon^2 n$ and at all times $t < \eta\epsilon n$ we have that $Y_t > -\frac{\eta^2}{3}\epsilon^2 n$ with probability tending to 1. As component sizes are excursion lengths of Y_t above its past minima, we conclude that by time $2(1 + \eta)\epsilon n$ we have explored completely at least one component of size at least $2(1 - \eta)\epsilon n$. As $N_t \leq n - t$ for $t > 2(1 - \eta)\epsilon n$ and $\eta < 1/4$ we have $\mathbf{E}[\eta_t - 1 \mid \mathcal{F}_{t-1}] \leq -\frac{\epsilon}{2}$. By optional stopping we immediately get that if $t_j > 2(1 - \eta)\epsilon n$ then $\mathbf{E}[t_{j+1} - t_j] \leq 2\epsilon^{-1}$. Thus if $\mathcal{C}$ is a component which we began discovering after time $2(1 - \eta)\epsilon n$ we have

$$\mathbf{P}(|\mathcal{C}| \geq \epsilon n) \leq \frac{2}{\epsilon^2 n}.$$

Denote by $X(\eta)$ the number of vertices of which $|C(v)| > \epsilon n$ which we began discovering after time $2(1 - \eta)\epsilon n$, then we learn that $\mathbf{E} X(\eta) \leq \frac{2}{\epsilon^2}$. Denote by $\mathcal{C}_1(\eta)$ the largest component which we began discovering after time $2(1 - \eta)\epsilon n$. Clearly if $|\mathcal{C}_1(\eta)| > \epsilon n$ then $X(\eta) > \epsilon n$, thus by Markov inequality

$$\mathbf{P}(|\mathcal{C}_1(\eta)| > \epsilon n) \leq \frac{2}{\epsilon^3 n} \rightarrow 0.$$

Thus we have proved that there exists a unique component of size between $2(1 - \eta)\epsilon n$ and $2(1 + \eta)\epsilon n$. Condition on this event and consider the graph remained on the complement of this component. This graph has m vertices where

$$|m - (n - 2\epsilon n)| < 2\eta\epsilon n,$$

and as $p = \frac{1+\epsilon}{n}$ we have that

$$\left| p - \left(1 - \frac{\epsilon}{m} \right) \right| \leq \frac{2\eta\epsilon + O(\epsilon^2)}{m}.$$

This graph is distributed as $G(m, p)$ restricted to the event that it does not contain a component of size between $2(1 - \eta)\epsilon n$ and $2(1 + \eta)\epsilon n$. By Theorem 1 we know that the event that there exists such a component has probability $o(1)$. Thus for any collection of graphs $\mathcal{B}$ on m vertices which does not contain such a component, the probability of $\mathcal{B}$ in the remaining graph is $(1 + o(1))\mathbf{P}_{m,p}(\mathcal{B})$ where $\mathbf{P}_{m,p}$ is the usual $G(m, p)$ probability measure. Thus, we conclude by Theorem 1 that for any integer $\ell > 1$ and $\eta' > 0$

$$\mathbf{P}\left(\left|\frac{|\mathcal{C}_\ell|}{2\epsilon^{-2}(n)\log(n\epsilon^3)} - 1\right| > \eta'\right) \rightarrow 0,$$

concluding the proof of the theorem. $\square$

Remark. With a little more effort it is possible to show for the super-critical case, that in the exploration process for any fixed ℓ , the ℓ -th largest component is discovered *after* the largest component is discovered.

Acknowledgments

The first author would like to thank Microsoft Research, in which parts of this research were conducted, for their kind hospitality.

The work was done partially while the authors were visiting the Institute for Mathematical Sciences, National University of Singapore in 2006. The visit was supported by the Institute.

References

- [1] Aldous D. (1997), Brownian excursions, critical random graphs and the multiplicative coalescent. *Ann. Probab.* **25**, 812–854.
- [2] N. Alon and J. H. Spencer, (2000) The probabilistic method, 2nd edition, Wiley, New York.
- [3] Bender E. A., Canfield E. R., McKay B. D. (1990), The asymptotic number of labeled connected graphs with a given number of vertices and edges. *Random Structures Algorithms*, **1**, 127–169.

- [4] Bollobás B. (1984), The evolution of random graphs. *Trans. Amer. Math. Soc.* **286**, 257–274.
- [5] Durrett, R. (1996), *Probability: Theory and Examples*, Second edition. Duxbury Press, Belmont, California.
- [6] Feller, W. (1971), *An introduction to probability theory and its applications. Vol. II.*, Second edition. John Wiley & Sons, Inc., New York-London-Sydney.
- [7] Karp, R. M. (1990), The transitive closure of a random digraph. *Random Structures Algorithms* **1**, 73–93.
- [8] Łuczak, T. (1990), Component behavior near the critical point of the random graph process. *Random Structures Algorithms* **1**, 287–310.
- [9] Łuczak T., Pittel B., Wierman J. C. (1994), The structure of a random graph at the point of the phase transition. *Trans. Amer. Math. Soc.*, **341**, 721–748.
- [10] Martin-Löf, A. (1986), Symmetric sampling procedures, general epidemic processes and their threshold limit theorems. *J. Appl. Probab.* **23**, 265–282.
- [11] Nachmias, A. and Peres, Y. (2005), The critical random graph, with martingales, submitted. Available at <http://www.arxiv.org/abs/math.PR/0512201>.
- [12] Nachmias, A. and Peres, Y. (2005) Critical percolation on random regular graphs, in preparation.
- [13] Spitzer, F. (1956), A combinatorial lemma and its application to probability theory. *Amer. Math. Soc.* **82**, 323–339.
- [14] Wright, E. M. (1977), The number of connected sparsely edged graphs. *Journal of Graph Theory* **1**, 317–330.

Asaf Nachmias: `asafnach(at)math.berkeley.edu`
 Department of Mathematics
 UC Berkeley
 Berkeley, CA 94720, USA.

Yuval Peres: `peres(at)stat.berkeley.edu`
Microsoft Research
One Microsoft way,
Redmond, WA 98052-6399, USA.